



## Semaglutide SL Suspension

**Chet Johnson Drug**  
204 Keller Ave N  
Amery, WI 54002  
**Phone:** 715-268-8121  
**Fax:** 715-268-4733

Date: \_\_\_\_\_

**Patient Name:** \_\_\_\_\_

**DOB:** \_\_\_\_\_

**Address:** \_\_\_\_\_

**Phone:** \_\_\_\_\_

**Prescriber Name:** \_\_\_\_\_

**Location:** \_\_\_\_\_

**Phone:** \_\_\_\_\_ **Fax:** \_\_\_\_\_

### Starting dose

Semaglutide 2 mg/mL SL suspension (Qty 18mL)

Place 0.5mL (1 mg) under the tongue, hold in place for 5 minutes then swallow once daily. Take on an empty stomach; Do not eat or drink for 30 minutes after dosing.

### 2 mg dose

Semaglutide 4 mg/mL SL suspension (Qty 18mL)

Place 0.5mL (2 mg) under the tongue, hold in place for 5 minutes then swallow once daily. Take on an empty stomach; Do not eat or drink for 30 minutes after dosing.

### 4 mg dose

Semaglutide 8 mg/mL SL suspension (Qty 18mL)

Place 0.5mL (4 mg) under the tongue, hold in place for 5 minutes then swallow once daily. Take on an empty stomach; Do not eat or drink for 30 minutes after dosing.

### 6 mg dose

Semaglutide 12 mg/mL SL suspension (Qty 18mL)

Place 0.5mL (6 mg) under the tongue, hold in place for 5 minutes then swallow once daily. Take on an empty stomach; Do not eat or drink for 30 minutes after dosing.

### 8 mg dose

Semaglutide 16 mg/mL SL suspension (Qty 18mL)

Place 0.5mL (8 mg) under the tongue, hold in place for 5 minutes then swallow once daily. Take on an empty stomach; Do not eat or drink for 30 minutes after dosing.

**Refills:** ☐ 1 Year ☐ 5 ☐ 3 ☐ 1 ☐ Zero

*When signed, pharmacy may increase dose based on patient's response unless noted under prescriber's signature.*

**Prescriber Signature:** **X** \_\_\_\_\_

**Date:** \_\_\_\_\_

**Notes:** \_\_\_\_\_

Information about semaglutide SL can be found at [www.subsema.com](http://www.subsema.com) or by scanning the QR code

